

**Project 3 of ST 755, Spring 2008**

**Due: Tuesday, 4/18/2008**

1. Consider the special case of the mixed model in project 2:

$$Y_{ij} = \beta + b_i + e_{ij} \quad i = 1, \dots, m, \quad j = 1, 2,$$

where  $b_i \sim N(0, \sigma_b^2)$  and  $e_{ij} \stackrel{\text{iid}}{\sim} N(0, \sigma_e^2)$ , independent of  $b_i$ . Suppose the true values of  $\sigma_b^2$  and  $\sigma_e^2$  are  $\sigma_b^2 = 0$  and  $\sigma_e^2 = 1$  (the true value of  $\beta$  is not very important for this problem). Denote  $\theta = (\beta, \sigma_b^2, \sigma_e^2)^T$  and let  $\theta_0$  denote its true value. Do the following:

- (a) Show that the Fisher information matrix  $I(\theta_0)$  for the MLE of  $\theta$  at  $\theta_0$  is equal to

$$I(\theta_0) = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

Note that  $I(\theta_0)$  is calculated at the true value  $\theta_0$ . Here  $m$  is the same as the  $n$  in Self and Liang (1987), and  $Y_i = (Y_{i1}, Y_{i2})^T$  is the same as  $X_i$ .

- (b) Show that the asymptotic distribution of the MLE of  $\beta$  will not be affected by the boundary issue. That is,  $\sqrt{m}(\hat{\beta} - \beta_0)$  will still have an asymptotic normal distribution with mean zero. Find the variance.
- (c) However, the asymptotic distribution of the MLEs of  $\sigma_b^2$  and  $\sigma_e^2$  will be affected by the boundary issue. That is,  $\sqrt{m}(\hat{\sigma}_b^2 - 0)$  and  $\sqrt{m}(\hat{\sigma}_e^2 - 1)$  will have asymptotic distributions with non-zero means (so the biases of  $\hat{\sigma}_b^2$  and  $\hat{\sigma}_e^2$  are at the order  $m^{-1/2}$ ). Find the means for these distributions. We know  $\hat{\sigma}_b^2$  will over-estimate its true value ( $= 0$ ). Will  $\hat{\sigma}_e^2$  also over-estimate its true value ( $= 1$ )?

2. Consider the following linear mixed model for longitudinal data

$$Y_{ij} = x_{ij}^T \beta + b_{i0} + b_{i1} t_{ij} + e_{ij} \quad i = 1, \dots, m, \quad j = 1, 2, \dots, n_j,$$

where  $\beta$  are fixed effects (of  $x$ ),  $t_{ij}$  are the times when  $Y_{ij}$  are measured,  $(b_{i0}, b_{i1})^T$  are random effects independent of the measurement errors  $e_{ij}$ . It is further assumed that

$$\begin{pmatrix} b_{i0} \\ b_{i1} \end{pmatrix} \sim N \left( \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} \sigma_{00} & \sigma_{01} \\ \sigma_{01} & \sigma_{11} \end{bmatrix} \right)$$

and  $e_{ij}$  are iid from  $N(0, \sigma_e^2)$ . We are interested in conducting the likelihood ratio test for the presence of the random effect  $b_{i1}$ .

- (a) Specify  $H_0$  and  $H_1$  in terms of model parameters for the above testing problem.
- (b) From the results presented in class, we know that under  $H_0$  the LRT statistic for testing the above hypothesis asymptotically has a distribution equal to the 50:50 mixture of  $\chi_1^2$  and  $\chi_2^2$ . Find out the critical values for the LRT for three levels 0.01, 0.05, 0.1;
- (c) If we used the regular LRT to test  $H_0$ , what is the consequence? Specifically, what are the actual sizes of the test if the nominal levels of the regular LRT are chosen to be 0.01, 0.05, 0.1?
- (d) Suppose we observed  $\text{LRT} = 2.45$ . What is the p-value?
- (e) Consider a special case of the above model:

$$Y_{ij} = \beta_0 + \beta_1 t_{ij} + b_{i0} + b_{i1} t_{ij} + e_{ij} \quad i = 1, \dots, 100, \quad j = 1, 2, \dots, 5,$$

where  $t_{ij} = j$ . Assume under  $H_0$  the model parameters take values  $\beta_0 = 2, \beta_1 = 1, \sigma_{00} = 1, \sigma_e^2 = 2$ . Use simulation to get the empirical distributions of the LRT statistic and p-value (use 1000 simulation runs). Compare this empirical distribution with its theoretical distribution. You can use the SAS procedure mixed to conduct the simulation.